

Compressibility of a two-dimensional extended Hubbard model

E. J. Calegari and S. G. Magalhães*

*Laboratório de Mecânica Estatística e Teoria da Matéria Condensada
Universidade Federal de Santa Maria, 97105-900 Santa Maria, RS, Brazil*

A. A. Gomes

*Centro Brasileiro de Pesquisas Físicas-CBPF
Rua Xavier Sigaud 150, 22290-180, Rio de Janeiro, RJ, Brazil*

February 2, 2008

Abstract

The compressibility of an extended $d-p$ Hubbard model is investigated by the Roth's two-pole approximation. Using the factorization procedure proposed by Beenen and Edwards, superconductivity with singlet $d_{x^2-y^2}$ -wave pairing is also considered. Within this framework, the effects of $d-p$ hybridization and Coulomb interaction U on the compressibility are studied carefully. It has been found that the compressibility diverges and then it becomes negative near the half-filling. Within Roth's method, it has been verified that an important contribution for the negative compressibility comes from the spin-correlation term $\langle S_j^+ S_i^- \rangle$ present in Roth's band shift. This correlation function plays an important role due to its high doping dependence. Also, its effects in the band shift and consequently in the compressibility are pronounced near the half-filling. The numerical results show that the hybridization acts in the sense of suppressing the negative compressibility near half-filling. Finally, the possibility of a connection between the negative compressibility and the phase separation is also discussed.

The non-Fermi liquid behavior observed in the normal state of the underdoped regime in cuprate systems has been subject of plenty study in recent years [1]. The existence of pseudogap, stripes and phase separation [1, 2] in the normal state, becomes appreciably hard the experimental and theoretical understanding of these systems. As the cuprates are strongly correlated electron systems, it is believed that the one-band Hubbard model can capture their main physical properties [2]. However, although this model gives interesting results, probably, a model which takes into account the hybridization between the d - and p -orbitals can be more adequate to treat these systems [3]. Recently, the present authors have applied the Roth's two-pole approximation [4] to study the effects of $d-p$ hybridization on the superconducting properties of an extended $d-p$ Hubbard model [3, 5, 6]. The authors followed the factorization procedure proposed by Beenen and Edwards [7] to consider d -wave superconductivity. However, near the half-filling, the obtained results show that the compressibility ($\kappa = \frac{dn_T}{d\mu}$) diverges in a critical value of total occupation number n_T (where $n_T = n_\sigma^d + n_{-\sigma}^d$), and then, it becomes negative. Nevertheless, the divergence may be associated with phase separation which occurs in the underdoped regime of the cuprate systems [2]. In this work, the nature of the divergence and the negative compressibility has been carefully investigated in the framework of the Roth's approximation [4].

*ggarcia@ccne.ufsm.br

The model used [3] is given by:

$$\begin{aligned}
H = & \sum_{i,\sigma} (\varepsilon_d - \mu) d_{i\sigma}^\dagger d_{i\sigma} + \sum_{i,j,\sigma} t_{ij}^d d_{i\sigma}^\dagger d_{j\sigma} + U \sum_i n_{i\uparrow}^d n_{i\downarrow}^d \\
& + \sum_{i,\sigma} (\varepsilon_p - \mu) p_{i\sigma}^\dagger p_{i\sigma} + \sum_{i,j,\sigma} t_{ij}^p p_{i\sigma}^\dagger p_{j\sigma} \\
& + \sum_{i,j,\sigma} t_{ij}^{pd} \left(d_{i\sigma}^\dagger p_{j\sigma} + p_{i\sigma}^\dagger d_{j\sigma} \right)
\end{aligned} \tag{1}$$

where μ is the chemical potential. The Green's functions necessary to treat the problem were obtained following the Roth's standart procedure [4], and they are found in reference [3]. The Roth's band shift $W_{\mathbf{k}\sigma} = W_{\mathbf{k}\sigma}^d + W_{\sigma}^{pd}$ is given in terms of correlation functions which can be evaluated from the obtained Green's functions (see Refs. [3, 6]). The $W_{\mathbf{k}\sigma}^d$ is given as:

$$n_{-\sigma}^d (1 - n_{-\sigma}^d) W_{\mathbf{k}\sigma}^d = h_{1\sigma} + \sum_{j \neq 0} t_{0j}^d e^{i\mathbf{k} \cdot \mathbf{R}_j} \bar{h}_{j\sigma} \tag{2}$$

where $\bar{h}_{j\sigma} = h_{2j\sigma} + h_{3j\sigma}$ and j stand for the nearest neighbors of the i site. The term $h_{1\sigma}$ is n_T -dependent but it is $\mathbf{k}$ -independent. The second term in the right side of Eq. (2) is both n_T and $\mathbf{k}$ -dependent. As discussed in Ref. [3], the term $h_{3j\sigma}$, which is directly associated with the superconductivity, is quite small and therefore, it can be disregarded. Finally, $h_{2j\sigma} = F_{j\sigma}^{(1)} + F_{j\sigma}^{(2)} + F_{3j\sigma}^{(3)}$ with,

$$F_{j\sigma}^{(1)} = - \frac{\alpha_{j\sigma} n_{0j\sigma}^d + \beta_{j\sigma} m_{j\sigma}}{1 - \beta_{\sigma} \beta_{-\sigma}} \tag{3}$$

$$F_{j\sigma}^{(2)} = \langle S_j^+ S_0^- \rangle = - \frac{\alpha_{j\sigma} n_{0j-\sigma}^d + \beta_{j\sigma} m_{j-\sigma}}{1 + \beta_{\sigma}} \tag{4}$$

$$F_{j\sigma}^{(3)} = - \frac{\alpha_{j\sigma} n_{0j-\sigma}^d + \beta_{j\sigma} (n_{0j-\sigma}^d - m_{j-\sigma})}{1 - \beta_{\sigma}} \tag{5}$$

where $n_{0j\sigma}^d$, $m_{j\sigma}$, $\alpha_{j\sigma}$ and $\beta_{j\sigma}$ are given in Ref. [3]. The correlation function $F_{j\sigma}^{(2)}$ is high doping δ dependent (where $\delta = 1 - n_T$), mainly near half-filling. Considering a model where $t_{0j}^d = t^d$ for the z neighbors, only one value ($j = 1$) of $h_{2j\sigma}$, named $h_{21\sigma}$, is need. The figures 1(a) and 1(b) show the behavior of W_{σ}^{pd} and $F_{1\sigma}^{(2)}$ as functions of n_T for different values of Coulomb interaction U and V_0^{pd} . The quantity V_0^{pd} stand for the $\mathbf{k}$ -independent hybridization [3]. It is clear from figures 1(a) and 1(b) that at least two important effects presented in the band shift $W_{\mathbf{k}\sigma}$ are responsible for the unexpected behavior of the compressibility. The first effect is associated with the spin-correlation term $\langle S_j^+ S_0^- \rangle$, which decreases significantly near half-filling as can be observed in figures 1(a) and 1(b). Therefore, due to the effect of $\langle S_j^+ S_0^- \rangle$, in the underdoping regime, the lower Hubbard band becomes very flat in the region of the point (π, π) [7], resulting in a peak in the density of states close to the charge-transfer gap (due to Coulomb interaction U). As consequence, below a critical doping δ_c , the chemical potential μ decreases. Moreover, the compressibility diverges in δ_c , and then, it becomes negative (see figs. 1(c) and 1(d)). Nevertheless, it has been verified that the hybridized shift W_{σ}^{pd} presented in $W_{\mathbf{k}\sigma}$ moves the divergence towards the half-filling when the hybridization is enhanced. This behavior is clear in figures 1(c) and 1(d), which show the chemical potential and the compressibility as functions of n_T . Using a systematic large- N expansion, in Ref. [8] the authors have studied the compressibility of the two-dimensional infinite- U one-band Hubbard model. They showed that a critical doping $\delta_c = 0.07 \pm 0.01$ exists, and it may sign the instability of the Fermi liquid phase (below δ_c) and the onset where phase separation can take place. Concerning the divergence in the compressibility, although the present model and procedure are distinct from those used in reference [8], the results are in agreement. Finally, it has been also verified that the $d-p$ hybridization moves the divergence in the compressibility towards the half-filling, mainly due to its effect on W_{σ}^{pd} . Therefore, the doping range where there are phase separation and Fermi liquid instability decreases if the hybridization is enhanced.




Figure 1: In (a), the hybridized shift W_{σ}^{pd} and the correlation function $F_{1\sigma}^{(2)}$ are shown as functions of n_T for $U = 8|t^d|$. The hybridizations $V = V_0^{pd}/|t^d|$ are given in the top of figure (a). (b) show W_{σ}^{pd} and $F_{1\sigma}^{(2)}$ for $V_0^{pd} = 1.0|t^d|$ and different Coulomb interactions. In (c), the chemical potential is shown as a function of n_T for $U = 8|t^d|$. The the hybridizations V are given in the top of figure (c). (d) dependence of the compressibility on n_T for the parameters identical to (c). ($k_B T = 0.0$ and $t^d = -0.5eV$).

References

- [1] J. L. Tallon and J. W. Loram, *Physica C*, **349** (2001) 53.
- [2] E. Dagotto, *Rev. Mod. Phys.*, **66** (1993) 763.
- [3] E.J. Calegari, S.G. Magalhães and A.A. Gomes, *Eur. Phys. J. B* **45** (2005) 485.
- [4] L.M. Roth, *Phys. Rev.* **184** (1969) 451.
- [5] E.J. Calegari, S.G. Magalhães and A.A. Gomes, *Intern. Journ. of Modern Phys. B*, **18** (2004) 241.
- [6] E.J. Calegari, S.G. Magalhães and A.A. Gomes, *Physica B* **359 - 361C** (2005) 560.
- [7] J. Beenen and D.M. Edwards, *Phys. Rev. B* **52** (1995) 13636.
- [8] Arti Tandon *et al.*, *Phys. Rev. Lett.* **83** (1999) 2046.